

Attached list 19b. Effective rate of acupuncture and sham acupuncture with other therapies.

Author	Disease	Acup schedule	Average daily dose of Acup	Total dose of Acup	Effect index	Groups	Number of participants	Before	After	Improvement rate
周君 (2010)	Spastic cerebral paralysis	Once a day for 10 days. Have one day break. Total 6 courses	0.91	54.55	MAS sore	Bobath, High pressure oxygen, Vojta (Sham, not inserted)	16	4.13	3.27	20.8%
						Bobath, High pressure oxygen, Vojta plus Acup	15	3.88	2.41	37.9%
夏隽晖 (2015)	Depression after stroke	Once a day for 6 day. Have one day off. Total 8 weeks	0.86	41.14	HAMD	Rehabilitation +Sham (not point, shallow insert)	48	29.9	19.9	33.4%
						Rehabilitation + Acup	50	30.5	13.5	55.7%
夏隽晖 (2015)b	Dysphagia after stroke	Once a day for 6 day. Have one day off. Total 8 weeks	0.86	41.14	SSA score	Rehabilitation +Sham (not point, shallow insert)	63	35.9	28.1	21.7%
						Rehabilitation + Acup	69	37.1	21.6	41.8%
Hawamdeh M (2015)	Chronic low back pain	3 times a week for 6 weeks	0.43	7.71		TENS + exercise	10	7	2	71%
						Sham TENS + exercise	10	7	5	29%
He D (2005)	chronic neck and shoulder pain	10 sessions during 3 to 4 weeks	0.41	4.08	VAS	E-Acup + aricular press	14	32	9	71.9%
						Sham Acup (inserted + ear acupress)	10	26	22	15.4%
Hopwood V	Stroke	3 sessions a week	0.43	5.14	Pain	Acup + physio + drugs	57	26.9	23.6	12%

(2008)		for 4 weeks			score at 6 week	Sham TENS + physio + drugs	48	22.1	20.1	9%
Sangdee C (2002)	Knee Osteoarthritis	3 times a week for 4 weeks	0.43	5.14	VAS	placebo tablet plus placebo EA (not inserted)	47	63.49	40.63	36.0%
						diclofenac tablet plus placebo EA	49	64.79	31.8	50.9%
						placebo tablet plus true EA (2 Hz)	48	66.87	18.63	72.1%
						diclofenac tablet plus true EA	49	57.63	22.04	61.8%
Man SC (2014)	post-stroke depression	3 sessions per week over 4 weeks	0.43	5.14	HAMD-17	DCEAS plus SSRI plus body electroacupuncture	23	25.8	14.2	45.0%
						sham (non-invasive cranial electroacupuncture, n-CEA) plus SSRI plus body electroacupuncture	20	28.1	16.9	39.9%
Yan T (2009) (Hong Kong)	acute stroke	5 days a week for 3 weeks	0.71	10.71	Up and Go scores after 3 weeks	TENS (with physiotherapy and occupational therapy)	19	67.5	30	55.6%
						Sham TENS (with physiotherapy and occupational therapy)	19	55.5	41.1	25.9%
						Conventional rehabilitation (with physiotherapy and occupational therapy)	18	46.4	55.4	-19.4%